

# Fungicidal Activities of Cu (II) Soaps Derived From Various Oils Treated at High Temperature for Biomedical Use

Bhutra R<sup>1</sup>, Sharma R<sup>2</sup> and Sharma A. K<sup>3\*</sup>

<sup>1</sup>Department of Chemistry, Rajesh Pilot Govt. Polytechnic College, Dausa, Raj, India

<sup>2</sup>Department of Chemistry, S.P.C. Govt. College, Ajmer, Raj, India

<sup>3</sup>Department of Chemistry, Govt. P.G. College, Jhalawar, Raj, India

\*Corresponding author: Sharma A. K, Department of Chemistry, Govt. P.G. College Jhalawar-326001 (Raj.) India, Tel: +91 9352669899, E-mail: sharmaarun423@gmail.com

**Citation:** Bhutra R, Sharma R, Sharma A. K (2018) Fungicidal Activities of Cu (II) Soaps Derived From Various Oils Treated at High Temperature for Biomedical Use. SAJ Biotechnol 5: 101

**Article history:** Received: 06 March 2018, Accepted: 29 March 2018, Published: 03 April 2018

## Abstract

Biologically potent compounds are one of the most important classes of materials for the upcoming generations. Increasing number of microbial infectious diseases and resistant pathogens create a demand and urgency to develop novel, potent, safe and improved variety of antimicrobial agents. This initiates a task for current chemistry to synthesize compounds that show promising activity as therapeutic agents with lower toxicity. Therefore, a substantial research is needed for their discovery and improvement. The antifungal activities of copper soaps derived from various oils treated at high temperatures have been evaluated by testing against *Alternaria alternata* and *Aspergillus niger* at different concentrations by agar plate technique and it is found that these Cu (II) soaps are very effective on these. The fungi toxicity results indicate that the strain of fungal species are susceptible towards these soaps and suggests that with the increase in concentration of copper soap it may increase further.

**Keywords:** Copper surfactants; Anti-fungal studies; Edible oils

## Introduction

Most of the antimicrobial agents are obviously organic compounds, which behave as good chelating agent. Their pathogenic activity is enhanced on complexing with various transition and toxic metals [1-2]. The biological effect of these derivatives depends on the nature and structure of ligands and their metal complexes and also on the presence of particular element. Many copper thiosemicarbazone complexes are found to have significant antitubercular, fungicidal and antitumor activities [3-5]. The use of copper linoleate as heavy duty wood preservative and many other biological activities of copper metal containing surfactants also have been studied [6-7]. Formulated proprietary brands of cuprous oxide are extensively employed as fungicides and seed dressings. Copper oxychloride has a number of applications, by far the most important being as an agricultural fungicide for which purpose it is extensively employed in formulated form as dusts, wettable powders and pastes [8].

Small quantities of copper soaps such as copper stearate, copper oleate and copper abietate (from resins), are employed mainly for rot-proofing textiles, ropes, etc. The main applications of copper sulfate are in the production of proprietary wood preservatives and agricultural fungicides [9-10]. In our earlier communication antifungal activities of neem, soyabean, sesame, groundnut and mustard oils, and their complexes with N,S donar ligands like Urea, thiourea, Benzothiazole have been reported. Mathur, *et al.* studied the biological evaluation and DNA binding cleavage studies of copper soaps and their complexes [11-19].

All the above studies suggest that the copper metal play a vital role in fungicidal activities. Various edible oils are widely used, easily available and economic. These facts led us to synthesis copper soaps of some oils and their treated oils remaining after frying, to study their micellar characteristics and role of copper metal in fungicidal activities for exploring their applications and other possible uses in various industries and agriculture. For this purpose, we have chosen some fungicidal activities of the synthesized copper soaps against easily available fungi *Alternaria alternata* and *Aspergillus niger*.

## Experimental

These oils are easily available in the Indian and chosen for the investigation. Their compositions are recorded in Table 1. All

the chemicals used were of LR/AR grade. Three samples for each oil have been prepared as fresh (untreated), treated oil at high temperature for 15 minutes and for 60 minutes.

Name of oil	Fatty Acids %					
	16:0	18:0	18:1	18:2	18:3	Other Acids
Groundnut Oil	10	4	61	18	-	C <sub>20</sub> -C <sub>24</sub> 7%
Mustard Oil	2	1	25	18	10	C <sub>20</sub> -C <sub>22</sub> 41%
Sesame Oil	8	4	45	41	-	-
Soyabean Oil	12	4	24	51	9	-

Table 1: Fatty Acid Composition of Oils Used For Copper Soap Synthesis

## Preparation of treated oils and soaps

About 500 g oil was taken in an iron pan. The oil was heated to the highest temperature and 5 g potato chips were deep fried while maintaining the frying temperature between 180-200 °C. The oil was heated for 15 min. in the open air and sunlight in iron pan at 180-200 °C. The volume of the oil was not replenished to the original volume after frying operations. The frying experiment with both oil samples at 180-200 °C for 60 minutes was conducted in similar manner.

## Preparation of copper soaps

Cu (II) soaps were prepared by refluxing the oils with 2N KOH solution and alcohol for about 3 h (Direct Metathesis). Copper soap so obtained was then washed with warm water and 10% alcohol at 50 °C and recrystallized using hot benzene. Molecular weight of copper soap was determined from saponification value of the oil sample prepared. The formation of copper soap was confirmed by using elemental analysis, UV, IR techniques.

**Fungicidal activities:** The general laboratory techniques followed in the course of this investigation are as suggested by Booth and Hawks worth [20-21].

## Sterilization of Glasswares

Glasswares used in our study were of pyrex brand. The glasswares viz. test tubes, conical flasks, pipettes (micro and macro), glass rods and petridishes were thoroughly washed after rinsing with chromic acid each time before sampling. The petriplates and other glassware were then sterilized in hot air oven at 160 °C for 24 hours before use.

## Culture Media Used

The cultured medium used for the growth of the organism in the present study was P.D.A. (Potato dextrose Agar). The following media was used in the present study

Potatoes-200g  
Dextrose-20g  
Agar-20g  
Distill water-1000ml

200g of potatoes were cleaned, cut into pieces and boiled in about 1000 ml of tap water for 2 h. Then, the contents were strained using muslin cloth. To this extract, 20 g of dextrose, 20 g of agar were added and made up to 1000 ml in graduated flask, before sterilizing the medium.

## Preparation of Sample Solutions

All the copper soaps derived from untreated and treated oils were tested for antifungal activity. The calculated amount of the soap was weighted in a standard flask and the solutions containing different concentration (1000 and 10000 ppm) of soaps in benzene were prepared

## Test Organism

In the present study, following two test organism are used:

- A. *Alternaria alternata*
- B. *Aspergillus niger*

The above organisms were isolated from their natural habitat and then purified, characterized and identified [22] (Figure 1a-1b).



Figure 1a: Presence of *Alternaria Alternata* on Tomato




Figure 1b: Presence of *Aspergillus Niger* on Bread

### Fungicidal Testing

1 ml of the soap solution was aseptically transferred into sterile petriplates. Into these plates, 20 ml of P.D.A. was poured and was mixed with soap solution by rotating the petriplates in clockwise and anticlockwise direction 3-4 times and was allowed to solidify.

After the solidification of the above medium, single hypha / spore of test organism were aseptically transferred in the centre of the petriplates. The plates were incubated at 30 °C for 3 days. After the period of incubation, the plates were observed for the growth of fungus in different concentration of the soap solution used in the present study. The data were statistically analysed according to the following formula:

$$\% \text{ Inhibition} = (C-T) \times 100 / C$$

Where C = diameter of fungal colony in control plates after 3 days

T = diameter of fungal colony in test plate after 3 days

### Results and Discussion

Copper soaps derived from untreated and treated oils have been screened for their antifungicidal activity against *Alternaria-alternata* and *Aspergillus-niger* at 1000 ppm and 10000 ppm by Agar-plate technique. Copper soaps showed moderate activities against both the fungi.

A perusal of Figure 2,3 and 4 reveals that all the copper soaps have significant fungitoxicity at 10000 ppm but their toxicity decreases markedly on dilution (at 1000 ppm). From the figs, it is apparent that their efficiency increases with their concentration. Thus it is evident that concentration plays a vital role in increasing the degree of inhibition. Fungicidal screening data revealed that at lower concentration the inhibition of growth is less as compared to higher concentration

From comparison of the results for both the fungi, it is found that all copper soaps are more potent (more toxic) against *Aspergillus niger* than against *Alternaria- alternata* i.e. inhibition of growth is higher for *Aspergillus niger* than inhibition of growth for *Alternaria alternata*. The results are concise with our earlier reported work on soaps with complexes derived from saturated fatty acids, edible and non-edible oils [23-26].

A perusal of the figs shows that CM is the least fungitoxic (% inhibition lowest) whereas CSo is the most toxic against both fungi. The activity (toxicity) of copper soaps derived from untreated oils are found to increase in the order




Figure 2: Antifungal activity of copper (ii) soaps derived from untreated oils




Figure 3: Antifungal activity of copper (ii) soaps derived from treated oils for 15 minutes

For copper soaps derived from treated oils for 15 minutes, the results are same as copper soaps derived from untreated oils. CM<sub>15</sub> is the least active and CSo<sub>15</sub> is the most active against both fungi. The order of activity of copper soaps derived from treated oils for 15 minutes is as follows [27]

$$CSo_{15} > CSe_{15} > CG_{15} > CM_{15}$$

For copper soaps derived from treated oils for 60 minutes, CM<sub>60</sub> is the least active but difference in activity of these copper soaps is not so high and order of activity is as follows:

$$CSo_{60} > CSe_{60} > CG_{60} > CM_{60}$$

From comparison of copper soaps derived from untreated and treated sample of oil, it is found that fungitoxicity increases with the increase of time of heating for oils i.e.




Figure 4: Antifungal activity of copper (ii) soaps derived from treated oils for 60 minutes

All the tests were performed in triplicate the standard deviation has been measured by the conventional measure of repeatability and the average was taken as final reading. The results of ANOVA for the antifungal activities for all soaps has been done, predicted  $R^2$  are in reasonable agreement and closer to 1.0. This confirms that the experimental data are well satisfactory [28]. The descriptive statistics results of Cu (II) soaps shown that SE and SD are 0.5 % which confirms satisfactory results in triplet. The result is statistically significant, by the standards of the study, due to  $p < F$  [29,30].

## Conclusion

It has been observed that the antifungal activity increases with the increase in the concentration of the solution. From comparison of copper soaps derived from untreated and treated sample of oil, it is found that fungitoxicity increases with the increase of time of heating for oils.

## Acknowledgement

The authors would like to thank Principal and Head of Department of Chemistry, S.D. Govt. College, BEAWAR and S.P.C. Govt. College, Ajmer- Raj. India for providing Laboratory Facilities.

## References

- Rashmi S, Arun KS (2017) Natural Edible Oils: Comparative Health Aspects Of Sesame, Coconut, Mustard (Rape Seed) and Groundnut (Peanut) A Biomedical Approach, Biomed J Sci Tech Res 1: BJSTR.MS.ID.000441.
- Sharma AK, Saxena M, Sharma R (2017) Acoustic Studies of Copper Soap- Urea complexes derived from Groundnut and Sesam oils. J Phy Studies 21: 4601-6.
- Tank P, Sharma AK, Sharma R (2017) Thermal Behaviour and Kinetics of Copper (II) Soaps and Complexes Derived from Mustard and Soyabean Oil. J Anal Pharm Res 4: 1-5.
- Sharma S, Sharma R, Sharma AK (2017) Synthesis, Characterization, and thermal degradation of Cu (II) Surfactants for sustainable green chem, Asian J Green Chem. 2: 129-140.
- Sharma AK, Saxena M, Sharma R (2018) Ultrasonic studies of Cu (II) Soaps derived from Groundnut and Sesame oils, Tenside. Surf Det 55: 127-134.

6. Bhati SK, Kumar A (2008) Synthesis of new substituted azetidinoyl and thiazolidinoyl-1,3,4-thiadiazino (6,5-b) indoles as promising anti-inflammatory agents, *Eur J Med Chem* 43: 2323-30.
7. Kumar A, Rajput CS (2009) Synthesis and anti-inflammatory activity of newer quinazoline-4-one derivatives. *Eur J Med Chem* 44: 83-90.
8. Mahajan K, Swami M, Singh RV (2009) Microwave synthesis, spectral studies, antimicrobial approach, and coordination behavior of antimony(III) and bismuth(III) compounds with benzothiazoline. *Russ J Coord Chem* 35: 179-80.
9. Sharma AK, Sharma S, Sharma R (2017) Thermal degradation of Cu (II) metallic Soaps and their Characterizations. A Pharmaceutical Application. *Chro Phar Sci* 1: 312-9.
10. Gunstone FD (1958) An introduction of the Chemistry of fats and fatty acids. Chapman & Hall Ltd., London, United Kingdom.
11. Khan S, Sharma R, Sharma AK (2017) Antifungal Activities of Copper Surfactants derived from Neem (*AzadiractaIndica*) and Karanj (*Pongamiapinnata*) Oils: A Pharmaceutical Application. *Glob J Pharmaceu Sci* 3: 1-6.
12. Tank P, Sharma R, Sharma AK (2017) A Pharmaceutical approach & Antifungal activities of Copper Soaps with their N & S donor complexes derived from Mustard and Soyabean oils, *Glob. J Pharmaceu Sci* 3: GJPPS.MS.ID.555619.
13. Sharma AK, Saxena M, Sharma R (2017) Synthesis, spectroscopic and fungicidal studies of Cu (II) soaps derived from groundnut and sesame oils and their urea complexes. *Bulletin of Pure and Applied Sciences*. 36: 26-37.
14. Saxena M, Sharma R, Sharma AK (2017) Micellar Features of Cu (II) Surfactants derived from Edible Oils. "ISBN 978-620-2-01906-4" LAP Lambert Academic Publishing Germany (2017).
15. Sharma AK, Singh N, Saxena M (2017) Synthesis, spectroscopic and biocidal activity of Cu (II) sesame complex. "ISBN 978-620-2-09609-6" LAP Lambert Academic Publishing Germany (2017).
16. Sharma AK, Saxena M, Sharma R (2018) Synthesis, Spectroscopic and Biocidal activities of environmentally safe Agrochemicals. *J Biochem Tech* 7: 1139-47.
17. Neha Mathur, Imran Ahmad, Ashok Kasana, Sonlata Bargotyia, Biplab Manna (2013) Biological Activities of Some New Environmentally Safe 2 Aminobenzothiazole Complexes of Copper (II) Derived Under Microwave Irradiation. *IAASCA* 5: 37-42.
18. Neha Mathur, Sonlata Bargotyia (2015) A facile synthesis and biological evaluation of some macrocyclic copper complexes. *IJPSR* 6: 2538-45.
19. Neha Mathur, Sonlata Bargotyia (2016) DNA binding and cleavage activities of macro cyclic metal complexes containing heteroatomic ligand. *Chem Sci Transactions* 5: 117-24.
20. Booth C (1971) "Methods in Microbiology". Acad Press N.Y. 4: 795.
21. Hawksworth OL (1974) 'Mycologists Hand Book' -An introduction to the principles of Taxonomy and nomenclature in the Fungi and Lichens. C. M.I. Kew, England.
22. Sharma R, Saxena M, Sharma N (2012) Synthesis, spectroscopic and fungicidal studies of copper soaps derived from mustard and soyabean oils and their urea complexes. *Int J Chem Sci* 10: 143-9.
23. Tank P, Sharma R, Sharma AK (2018) Viscometric Studies of Cu (II) surfactants derived from mustard oil in benzene at 303.15K. *Tenside Surf Det.*
24. Bhutra R, Sharma R, Sharma AK (2017) Viscometric and CMC studies of Cu(II) surfactants derived from untreated and treated groundnut and mustard oils in non-aqueous solvent at 298.15 K. *J Inst Chemists* 90: 29-47.
25. Sharma S, Sharma R, Heda LC, Sharma AK (2017) Kinetic parameters and Photo Degradation studies of Copper Soap derived from Soybean Oil using ZnO as a Photo catalyst in Solid and Solution Phase. *J Inst Chemists* 89: 119-36.
26. Khan S, Sharma R, Sharma AK (2018) Acoustic studies and other Acoustic Parameters of Cu(II) Soap derived from nonedible Neem oil (*Azadiracta indica*), in Non-aqueous media at 298.15. *Acta Ac united Ac.* 104: 277-83.
27. Joram A, Sharma R, Sharma AK (2018) Thermal degradation of complexes derived from Cu (II) groundnut soap (*Arachis hypogaea*) and Cu (II) sesame soap (*Sesamum indicum*). *Int J Res in Phys Chem and Chem Phys*.
28. Garg BS, Kumar DN (2003) Spectral studies of complexes of nickel (II) with tetradentate schiff bases having N2O2 donor groups, *Spectrochim. Acta* 59A: 229-34.
29. Bhutra R, Sharma R, Sharma AK (2018) Synthesis, Characterization and fungicidal activities of Cu (II) surfactants derived from groundnut and mustard oils treated at high temperatures. *J Inst Chemists* 90: 66-80.
30. Sharma AK, Sharma R, Saxena M (2018) Biomedical and antifungal application of Cu(II) soaps and its urea complexes derived from various oils, Open access *J Trans Med Res* 2: 40-3.